













































- [11] L. R. M. MAAS, D. BENIELLI, J. SOMMERIA, AND F.-P. A. LAM, *Observation of an internal wave attractor in a confined stable stratified fluid*, *Nature*, 388 (1997), pp. 557–561.
- [12] P. MÜLLER AND X. LIU, *Scattering of Internal Waves at Finite Topography in Two Dimensions. Part I: Theory and Case Studies*, *J. Phys. Oceanogr.*, 30 (2000), pp. 532–549.
- [13] ———, *Scattering of Internal Waves at Finite Topography in Two Dimensions. Part II: Spectral Calculations and Boundary Mixing*, *J. Phys. Oceanogr.*, 30 (2000), pp. 550–563.
- [14] F. PETRELIS, S. LLEWELLYN SMITH, AND W. R. YOUNG, *Tidal Conversion at a Submarine Ridge*, *J. Phys. Oceanogr.*, 36 (2006), pp. 1053–1071.
- [15] L. RAINVILLE AND R. PINKEL, *Propagation of Low-Mode Internal Waves through the Ocean*, *J. Phys. Oceanogr.*, 36 (2006), pp. 1220–1236.
- [16] H. SANDSTROM, *On topographic generation and coupling of internal waves*, *Geophys. Astrophys. Fluid Dyn.*, 7 (1975), pp. 231–270.
- [17] L. ST. LAURENT AND C. GARRETT, *The Role of Internal Tides in Mixing the Deep Ocean*, *J. Phys. Oceanogr.*, 32 (2002), pp. 2882–2899.